

Neoplasia of the Testis Histopathology Reporting Guide Retroperitoneal Lymphadenectomy



Main' You's	Ketropei	itoneai L	lymphadenectomy
Family/Last name			Date of birth DD - MM - YYYY
Given name(s)			
Patient identifiers		D	ate of request Accession/Laboratory number
			DD - MM - YYYY
Elements in black text ar	_	ey text are NON	SCOLE OF THIS DATASET
CLINICAL INFORMATI	ON I		SPECIMEN(S) SUBMITTED continued
☐ Information not provided ☐ Information provided (select all that apply) ☐ Previous history of testicular cancer, specify			Brain Lung Liver Other, specify
Previous therap	y, specify		
Other clinical in	formation, <i>specify</i>		SIZE OF LARGEST LYMPH NODE Cannot be assessed
	ionnacion, specin,		Maximum dimension
			mm
			Additional dimensions
SERUM TUMOUR MARK	ERS		mm x mm
Not provided			
Specify serum tum	markers within normal lii our markers used, level a in (select all that apply)		SIZE OF LARGEST NODAL METASTASIS Cannot be assessed
Date	□AFP	ug/L	Maximum dimension
LDH	□ b-HcG	IU/L	Additional dimensions
CDECIMEN(C) CUDMIT	TED (calcat all that apply)	- Ma-1	mm x mm
Not specified Retroperitoneal lym	phadenectomy, <i>specify n</i>		BLOCK IDENTIFICATION KEY (List overleaf or separately with an indication of the nature and origin of all tissue blocks)
	→ ○	No disease Necrosis Viable tumour	HISTOLOGICAL TUMOUR TYPE (select all that apply) (Value list based on the World Health Organization Classification of Urinary and Male Genital Tumours (2022))
	→ ○	No disease Necrosis Viable tumour	✓ Viable tumour ✓ Absent ✓ Present
	' U	No disease Necrosis Viable tumour	%

	n cell tumour <i>, spec</i>	cify type and percer	itage
		-	
		→	
		→	
Othe	er, <i>specify</i>		
	, , -, ,		
GIN	STATUS		
	not be assessed		
	involved		
	stance of tumour fr argin	om closest	mm
Sp	ecify closest margi	n(s), if possible	
Invo	lved		
Sp	ecify margin(s), if	possible	
RANG	DDAL EXTENSIO	N MA	
	DDAL EXTENSION	N 🗐	
Inde	terminate	N	
Inde	terminate identified	N D	
Inde Not	terminate identified	N	
Inde Not Pres	terminate identified ent		ition) ^a
Inde Not Pres	terminate identified ent	(UICC TNM 8 th ed	ition) ^a
Inde Not Pres	terminate identified ent GICAL STAGING	(UICC TNM 8 th ed	ition) ^a
Inde Not Pres HOLO M De	terminate identified ent GICAL STAGING scriptor (only if appost-therapy	(UICC TNM 8 th ed	ition) ^a
Inde Not Pres HOLO M De y -	terminate identified ent GICAL STAGING scriptor (only if ap post-therapy al lymph nodes (p	(UICC TNM 8 th edopplicable)	
Inde Not Pres HOLO M De y - giona NX ^b	terminate identified ent GICAL STAGING scriptor (only if appost-therapy al lymph nodes (p	(UICC TNM 8 th edopplicable) DN) odes cannot be ass	
Inde Not Pres HOLO M De y - giona NX ^b NO	terminate identified ent GICAL STAGING scriptor (only if appost-therapy al lymph nodes (p Regional lymph n	(UICC TNM 8 th edopplicable)	essed
Inde Not Pres HOLO M De y - giona NX ^b NO	terminate identified ent GICAL STAGING scriptor (only if appost-therapy al lymph nodes (p Regional lymph n No regional lymph Metastasis with a greatest dimensic	(UICC TNM 8 th edopplicable) DN) odes cannot be asson node metastasis lymph node mass in and five or fewer	essed 2 cm or lespositive no
Inde Not Pres HOLO M De	dentified ent GICAL STAGING Scriptor (only if appost-therapy Al lymph nodes (p Regional lymph n No regional lymph Metastasis with a greatest dimension none more than 2 Metastasis with a	(UICC TNM 8 th ed oplicable) ON) odes cannot be ass in node metastasis lymph node mass in and five or fewer or cm in greatest din lymph node mass	essed 2 cm or lest positive nonension more than
Inde Not Pres HOLC M De y - NX ^b NO N1	dentified ent GICAL STAGING Scriptor (only if appost-therapy Al lymph nodes (p Regional lymph n No regional lymph Metastasis with a greatest dimension none more than 2 Metastasis with a 2 cm but not more	(UICC TNM 8 th ed oplicable) ON) odes cannot be ass in node metastasis lymph node mass in and five or fewer or and five or fewer or cm in greatest din lymph node mass re than 5 cm in greatest	essed 2 cm or lest positive nonension more than atest
Inde Not Press IOLC M De y - giona NX ^b N0 N1	dentified ent DGICAL STAGING Scriptor (only if appost-therapy Allymph nodes (programmer) Regional lymph nodes destates with a greatest dimension none more than 2 cm but not more dimension; or monone more than 5	(UICC TNM 8 th ed oplicable) ON) odes cannot be asson node metastasis lymph node masson and five or fewer or and five or fewer or greatest din lymph node masson than 5 cm in greater than 5 cm; or evidence of com; or evidence of	essed 2 cm or lest positive notes than atest positive,
Inde Not Press IOLC 4 De y - giona NX ^b N0 N1	dentified ent DGICAL STAGING Scriptor (only if appost-therapy Allymph nodes (page 1 lymph nodes (page 2 lymph nodes 3 lymph nodes 4 lymph nodes 3 lymph nodes 3 lymph nodes 3 lymph nodes 4 lymph no	(UICC TNM 8 th ed oplicable) ON) odes cannot be asson node metastasis lymph node masson and five or fewer or and five or fewer or greatest din lymph node masson than 5 cm in greater than 5 cm; or evidence of com; or evidence of	essed 2 cm or lest positive nonension more than atest positive, f extranod

	Distant metastasis (pM) (if resected)					
	No distant metastasesM1 Distant metastasisM1a Non-regional lymph node(s) or lungM1b Other sites					
а	Reproduced with permission. Source: UICC TNM Classification of Malignant Tumours, 8th Edition, eds by James D. Brierley, Mary K. Gospodarowicz, Christian Wittekind. 2016, Publisher Wiley (incorporating any errata published up until 12th July 2024).					
)	NX should be used only if absolutely necessary.					